

Complexation and redox interactions between aqueous plutonium and manganese oxide interfaces

Dawn A. Shaughnessy¹, Heino Nitsche^{1,2}, R. Jeffrey Serne³, David K. Shuh¹, Glenn A. Waychunas⁴, Corwin H. Booth¹, Kirk J. Cantrell³, Richard E. Wilson^{1,2}

¹Chemical Sciences Division, Glenn T. Seaborg Center, Lawrence Berkeley National Laboratory
MS 70A-1150, Berkeley, CA 94720, USA

dakeeney@lbl.gov

²Department of Chemistry, University of California
Berkeley, CA 94720, USA

³Pacific Northwest National Laboratory
Richland, WA 99352, USA

⁴Earth Sciences Division, Lawrence Berkeley National Laboratory
MS 90-1116, Berkeley, CA 94720, USA

Manganese oxides are present throughout the environment, and their interaction with plutonium is of interest for the remediation of nuclear waste. There can be several manganese valence states in one oxide mineral, and plutonium can simultaneously exist in up to four different oxidation states. Therefore, there is a high probability that redox reactions will occur between the manganese and plutonium ions sorbed to the oxide surface.

To understand these interfacial reactions, we are studying the sorption of aqueous plutonium ions in single oxidation states on oxide minerals as a function of pH, ionic strength, and concentration. The sorbed plutonium complexes have been analyzed spectroscopically to determine oxidation states and the structures of the sorbed species.

Preferred session number: 3. (Solution Chemistry)

Last name: Shaughnessy

First name: Dawn

Title: Dr.

Mailing Address: Lawrence Berkeley National Laboratory, 1 Cyclotron Road, MS 70A-1150,
Berkeley, CA 94720, USA

TEL: +1-510-486-7307

FAX: +1-510-486-7444

e-mail: dakeeney@lbl.gov

Number of accompanied persons who will not attend the technical sessions: 0 or 1 (to be determined)